

## Anti-h CA125 4601 SPTN-5

### Product overview

---

<b>Catalog number</b>	100598
<b>Specificity</b>	Antibody recognizes human cancer antigen 125 (CA125)
<b>Description</b>	Monoclonal mouse antibody, cultured <i>in vitro</i> under conditions free from animal-derived components.
<b>Product buffer solution</b>	50 mM Na-citrate, pH 6.0, 0.9 % NaCl, 0.095 % NaN <sub>3</sub> as a preservative
<b>Shelf life and storage</b>	36 months from manufacturing at 2–8 °C
<b>Subclass</b>	IgG <sub>1</sub>
<b>Analyte description</b>	Cancer antigen 125 is a peptide epitope on the mucin glycoprotein MUC16. CA125 is the most widely used serum biomarker to monitor epithelial ovarian cancer. It is also used for the differential diagnosis of pelvic masses.

### Parameters tested on each lot

---

<b>Product appearance</b>	Liquid, may turn slightly opaque during storage
<b>Product concentration</b>	5.0 mg/ml (+/- 10 %)
<b>Immunoreactivity</b>	80–120 % compared to the reference sample in an FIA test
<b>IEF Profile</b>	6.2–7.1
<b>Purity</b>	≥ 95 %

### Kinetic parameters

---

<b>Association rate constant</b>	Not Determined (N/D)
<b>Dissociation rate constant</b>	N/D
<b>Affinity constant</b>	N/D
<b>Determination method</b>	-
<b>Determination antigen</b>	-



#### Legal disclaimer

**Cross-reactivities** Does not recognize carbohydrate antigen 19-9 (CA19-9), cancer antigen 15-3 (CA15-3), CYFRA 21-1, carcinoembryonic antigen (CEA),  $\alpha$ -fetoprotein (AFP), or human epididymis protein 4 (HE4).

**Epitope** Antibody 4601 is OC125-like and belongs to group A (ISOBM classification) as described in Nustad et al. 1996.

**Pair recommendations**

		DETECTION	
		4601	4602
CAPTURE	4601	-	+
	4602	+	-

Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations are only indicative.

**Platforms tested** FIA

**Antigens tested** Native CA125 antigen, Lee Biosolutions 151-25.

TEMPERATURE, TIME	RESULT
-70 °C, 21 days	OK
-20 °C, 21 days	OK
+4 °C, 21 days	OK
+35 °C, 21 days	OK
+45 °C, 7 days	OK

Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. Please note that the shelf life given on the first page is based on real time stability testing at 2–8 °C in the product buffer.

**Miscellaneous** -

**References** Nustad K, Bast RC, O'Brien TJ et al. (1996) Specificity and affinity of 26 monoclonal antibodies against the CA 125 antigen: First report from the ISOBM TD-1 Workshop. Tumor Biol 17:196-219.



**Legal disclaimer**