

## Anti-h CA15-3 4404 SPTN-5

### Product overview

---

<b>Catalog number</b>	100584
<b>Specificity</b>	Antibody recognizes MUC-1 core protein of cancer antigen 15-3
<b>Description</b>	Monoclonal mouse antibody, cultured <i>in vitro</i> under conditions free from animal-derived components.
<b>Product buffer solution</b>	50 mM Na-citrate, pH 6.0, 0.9 % NaCl, 0.095 % NaN <sub>3</sub> as a preservative
<b>Shelf life and storage</b>	24 months from manufacturing at 2–8 °C
<b>Subclass</b>	IgG <sub>1</sub> , λ light chain
<b>Analyte description</b>	Cancer Antigen 15-3 is defined by the use of two monoclonal antibodies, one specific for MUC-1 protein core and another antibody specific for a sialylated carbohydrate epitope on the MUC-1 protein. CA15-3 is a widely used serum marker in breast cancer patients for monitoring their response to cancer therapy.

### Parameters tested on each lot

---

<b>Product appearance</b>	Liquid, may turn slightly opaque during storage
<b>Product concentration</b>	5.0 mg/ml (+/-10 %)
<b>Immunoreactivity</b>	80–120 % compared to the reference sample in an FIA test
<b>IEF Profile</b>	5.9–7.1
<b>Purity</b>	≥ 95 %

### Kinetic parameters

---

<b>Association rate constant</b>	Not Determined (N/D)
<b>Dissociation rate constant</b>	N/D
<b>Affinity constant</b>	N/D
<b>Determination method</b>	-
<b>Determination antigen</b>	-



#### Legal disclaimer

**Cross-reactivities** N/D

**Epitope** Antibody binds to tandem repeat APDTRPAPGSTAPPAHGVTs of the MUC-1 core protein.

**Pair recommendations**

		DETECTION	
		4404	R4406
CAPTURE	4404	-	+
	R4406	+	-

Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations are only indicative.

**Platforms tested** FIA

**Antigens tested** Native CA15-3 antigen, Medix Biochemica 151-53A.

Product stability	TEMPERATURE, TIME	RESULT
	-70 °C, 21 days	OK
	-20 °C, 21 days	OK
	+4 °C, 21 days	OK
	+35 °C, 21 days	OK
	+45 °C, 7 days	OK

Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. Please note that the shelf life given on the first page is based on real time stability testing at 2–8 °C in the product buffer.

**Miscellaneous** -

**References** -



**Legal disclaimer**